

SUTAB[®]

(sodium sulfate, magnesium sulfate, and potassium chloride)

Tablets

1.479 g/0.225 g/0.188 g

Your procedure will be performed by Doctor: _____

Address: _____

Date of procedure: _____ Arrive at: _____ AM/PM

Comments: _____

How to Prep for Your Upcoming Colonoscopy

On the Day Before Your Procedure

What You CAN Do

- You may have clear liquids

What You CANNOT Do

- Do not drink milk or eat or drink anything colored red or purple
- Do not drink alcohol
- Do not take other laxatives while taking SUTAB
- Do not take oral medications within 1 hour of starting each dose of SUTAB
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin, chlorpromazine, or penicillamine, take these medications at least 2 hours before and not less than 6 hours after administration of each dose of SUTAB.

Liquids That Are OK to Drink


- Coffee or tea (no cream or nondairy creamer)
- Fruit juices (without pulp)
- Gelatin desserts (no fruit or topping)
- Water
- Chicken broth
- Clear soda (such as ginger ale)



DOSE 1 – On the Day Prior to Colonoscopy

STEP 1 Open 1 bottle of 12 tablets.

STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water, and drink the entire amount of water over 15 to 20 minutes.



IMPORTANT: If you experience preparation-related symptoms (for example, nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.

Drink additional water

STEP 3 Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

STEP 4 Approximately 30 minutes after finishing the second container of water, fill the provided container with 8 ounces of water (halfway to the fill line), and drink the entire amount over 30 minutes.

DOSE 2 – Day of the Colonoscopy

- Continue to consume only clear liquids until after the colonoscopy
- The morning of the colonoscopy (5 to 8 hours prior to the colonoscopy and no sooner than 4 hours from starting Dose 1), open the second bottle of 12 tablets
- Repeat STEP 1 to STEP 4 from Dose 1 (**IMPORTANT: You must complete all SUTAB tablets and required water at least 2 hours before colonoscopy.**)